

References:

- Mayr, E. 1935. Birds collected during the Whitney South Sea Expedition, XXX: descriptions of twenty-five new species and subspecies. *Amer. Mus. Novit.* No. 820: 1-6.
 Salomonsen, F. 1962. Whitehead's Swiftlet (*Collocalia whiteheadi* Ogilvie-Grant) in New Guinea and Melanesia. *Vidensk. Medd. fra Dansk Naturh. Foren.* 125: 509-512.
 Somadikarta, S. 1967. A recharacterization of *Collocalia papuensis* Rand, the Three-toed Swiftlet. *Proc. U.S. Nat. Mus.* 124 (3629): 1-8.
 Address. Dr. S. Dillon Ripley, Smithsonian Institution, Washington, D.C. 20560, U.S.A.
 © British Ornithologists' Club 1983

The distribution of the races of *Arremon aurantirostris* (Emberizinae) in Panama

by Storrs L. Olson

Received 27 October 1982

Although 3 subspecies of the Orange-billed Sparrow *Arremon aurantirostris* are recognized in Panama, the characters and distribution of 2 of these have been inaccurately represented in the literature. The dark race *A. a. rufidorsalis* Cassin, 1865 (type locality, Turrialba, Costa Rica), of the Caribbean slope of Middle America extends into northwestern Panama in Bocas del Toro and requires no further comment. The nominate subspecies *A. a. aurantirostris* Lafresnaye, 1847 (type locality "Panama"), is usually stated to range from the Pacific slope of Costa Rica east in Panama to the former Canal Zone (e.g. Hellmayr 1938, Paynter 1970). A third subspecies, *A. a. strictocollaris* Todd, 1922 (type locality, "Sautata" = Saotata, lower Río Atrato, Chocó, Colombia), has been considered to range from extreme eastern Panama into the adjacent parts of Chocó, in northwestern Colombia (Hellmayr 1938, Paynter 1970). These ranges imply a gap in the distribution of the species that does not in fact exist. Furthermore, although Todd (1922) characterized *A. a. strictocollaris* as having a more yellowish green dorsum, a narrower black pectoral band, and lighter underparts than *A. a. aurantirostris*, none of these characters actually holds.

Chapman (1925) and Hellmayr (1938) incorrectly regarded *A. a. strictocollaris* as being intermediate between *A. a. aurantirostris* and the South American subspecies, such as *A. a. occidentalis* and *A. a. erythrorhynchus*, which have a narrower black pectoral band. In the series of specimens from northwestern Colombia and from most of Panama that I have examined, the pectoral band was quite variable in width and it may also be considerably affected by the manner of preparation of the skin. The pectoral band in the holotype of *A. a. strictocollaris*, which I examined at the Carnegie Museum of Natural History, falls within the range of variation of *A. a. aurantirostris* and thus does not approach the South American subspecies in this respect. In addition (*contra* Todd 1922), the dorsum of *A. a. strictocollaris* is *not* more yellowish green than in *A. a. aurantirostris*—in fact, the opposite is true—nor are the purported differences in colour of the underparts of the 2 subspecies apparent to me.

Thus, although weakly differentiated from *A. a. aurantirostris*, *A. a. strictocollaris* may nevertheless be distinguished by (1) the decidedly orange-yellow rather than yellow coloration of the bend of the wing and (2) the slightly darker, more olivaceous dorsum. The broad-banded forms *A. a.*

strictocollaris and *A. a. aurantiirostris* differ from one another in almost exactly the same manner as the narrow-banded forms *A. a. occidentalis* and *A. a. erythrorhynchus* differ from each other.

Individuals with orange-yellow bends of the wing occur not only in the range traditionally ascribed to *A. a. strictocollaris* but also west to the area of the Canal Zone, where both *strictocollaris* and *aurantiirrostris* occur. Thus it becomes of importance to determine the identity of the specimens upon which Lafresnaye (1847) based the name *Arremon aurantiirrostris*, as these would presumably have come from the central part of Panama where either subspecies might be represented.

Although Stone (1899) once regarded a specimen in the collections of the Academy of Natural Sciences of Philadelphia (ANSP 7789) to be *the* type of Lafresnaye's name, Bangs (1930) showed that a specimen in the Museum of Comparative Zoology (MCZ 76675) had an equal claim and that both would have to be considered co-types. I was able to compare the Philadelphia specimen with the series in the National Museum of Natural History, Smithsonian Institution (USNM), and found that, although it is somewhat faded, it definitely has an orange-yellow bend of the wing. I then sent a series of fresh specimens of both the subspecies in question to R. A. Paynter, Jr., who compared them with the co-type of *aurantiirrostris* in the MCZ. He reported that "the dorsum is hopeless, as the bird is ancient, but the bend of the wing is definitely *yellow*, not *orangish*" (Paynter *in litt.* 24 August 1982). Accepting this, I designate MCZ 76675 as the lectotype of *Arremon aurantiirrostris* Lafresnaye, 1847, in order to preserve the current nomenclature, as the Philadelphia co-type would have to be referred to the population that now bears the name *strictocollaris*.

The characters and distribution of the two subspecies of Orange-billed Sparrow that occupy most of Panama may be summarized as follows:—

Arremon aurantiirrostris aurantiirrostris Lafresnaye, 1847. Dorsum yellowish green, bend of wing yellow. Pacific slope of Costa Rica and Panama east to the Canal Zone, where found on both slopes, and extending as far west along the Caribbean side as the Atlantic drainage of Coclé (El Uracillo). Mixes with *A. a. strictocollaris* in the Canal Zone area and along the Caribbean coast probably as far east as western San Blas (2 of 3 specimens from Mandinga appear to be more similar to the nominate form).

Arremon aurantiirrostris strictocollaris Todd, 1922. Similar to *A. a. aurantiirrostris* but dorsum darker, more olivaceous, bend of wing more deeply coloured, orange-yellow rather than yellow. Vicinity of Canal Zone, eastern Panama Province and eastern Colón, eastward along both slopes to north-western Colombia in Chocó.

Specimens (USNM) from Panama and Colombia examined.

A. a. rufidorsalis: PANAMA. BOCAS DEL TORO: Changuinola River (2); Almirante (2).

A. a. aurantiirrostris: PANAMA. CHIRIQUI: Divalá (3); San Felix (1); El Volcan, Palo Santo (1); Las Lajas (1); Puerto Armuelles (1). VERAGUAS: Soná (5); Santa Fé (1); "Veragua" (1). LOS SANTOS: Cerro Hoya (1). COCLE: El Copé (1); El Uracillo (2). CANAL ZONE: Río Indio (2); Lion Hill (3); Chiva Chiva (1); Gatun (1); Summit (1); Curundu (1). PANAMA PROVINCE: Peluca Hydrographic Station (1). COLON: Chilar (1). SAN BLAS: Mandinga (2).

A. a. strictocollaris: PANAMA. CANAL ZONE: Lion Hill (2); Tabernilla (2); Gatun (2). PANAMA PROVINCE: Utivé (1); Cerro Chucantí (3); Cerro Azul (1). COLON: Porto Bello (2). SAN BLAS: Mandinga (1); Armila (1). DARIEN: Río Jaqué (8); Tacarcuna Village (9); La Laguna (1); Cerro Pirre (1); Cana (2). COLOMBIA. CHOCO: Acandí (5).

Acknowledgements. I am grateful to Frank B. Gill for transporting the Philadelphia co-type of *Arremon aurantirostris* to Washington for my examination, to Raymond A. Paynter, Jr. for his remarks on the Harvard co-type and to Kenneth C. Parkes for access to the collections in the Carnegie Museum of Natural History and for comments on the manuscript. This is contribution Number 13 of the Wetmore Papers, a project supported in part by trust funds from the Smithsonian Institution for completing unfinished work and study of undescribed material left by the late Alexander Wetmore.

References:

- Bangs, O. 1930. Types of birds now in the Museum of Comparative Zoölogy. *Bull. Mus. Comp. Zool.* 70: 147-426.
- Chapman, F. M. 1925. Descriptions of new birds from Colombia, Ecuador, Peru, and Argentina. *Amer. Mus. Novit.* 160: 1-14.
- Hellmayr, C. E. 1938. Catalogue of birds of the Americas. Part XI. *Zool. Ser. Field Mus. Nat. Hist.* 13 (11): vi + 662 pp.
- Paynter, R. A., Jr. 1970. Subfamily Emberizinae, pp. 3-214 in *Check-list of Birds of the World*. Vol. XIII (Paynter, R. A., Jr. & Storer, R. W.—Eds). Cambridge, Mass.: Museum of Comparative Zoology.
- Stone, W. 1899. A study of the type specimens of birds in the collection of the Academy of Natural Sciences of Philadelphia, with a brief history of the collection. *Proc. Acad. Nat. Sci. Philadelphia* 1899 (part 1): 5-62.
- Todd, W. E. C. 1922. New forms of finches and tanagers from tropical America. *Proc. Biol. Soc. Washington* 35: 89-94.
- Address.* S. L. Olson, National Museum of Natural History, Smithsonian Institution, Washington, D.C. 20560, U.S.A.

© British Ornithologists' Club 1983

Notes on the extinct *Argusianus bipunctatus* (Wood)

by G. W. H. Davison

Received 2 November 1982

All published sources, including Peters (1934), Delacour (1951) and Warren (1966), cite incorrectly the type description of *Argusianus bipunctatus* (Wood).

The description accepted as the first by these writers was a letter written by T. W. Wood on 22 June 1871 (the author's manuscript date) and published presumably later than June (Wood 1871a). This letter gives an engraving of the type specimen, a description, and the proposed name *Argus bipunctatus* attached both to the illustration (which is on an earlier page) and to the description. However, this letter states that "a letter of mine appeared in the 'Field' newspaper of April 8th. ult. . .".

Reference to *The Field* for that date shows that Wood (1871b) had published the same engraving, with a nearly identical text, and proposed the name *bipunctatus*, approximately 3 months earlier than his June letter. Correct citation of the type description should therefore be:

Argus bipunctatus Wood, *The Field*, 8 April 1871, p. 281.

Subsequent mention of this species in publications has included no more details than were given by Wood himself. The type and only known